This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.

This Page Blank (uspto)

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6:

C07D 417/12, A61K 31/425

(11) International Publication Number:

WO 99/31094

A1

(43) International Publication Date:

24 June 1999 (24.06.99)

(21) International Application Number:

PCT/EP98/08154

(22) International Filing Date:

14 December 1998 (14.12.98)

(30) Priority Data:

9726568.0

16 December 1997 (16.12.97) GB

(71) Applicant (for all designated States except US): SMITHKLINE BEECHAM PLC [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB).

(72) Inventors; and

- (75) Inventors/Applicants (for US only): BLACKLER, Paul, David, James [GB/GB]; SmithKline Beecham Pharmaceuticals, Old Powder Mills, Near Leigh, Tonbridge, Kent TN11 9AN (GB). LEE, David, C. [GB/GB]; SmithKline Beecham Pharmaceuticals, The Frythe, Welwyn, Hertfordshire AL6 9AR (GB). SASSE, Michael, John [GB/GB]; SmithKline Beecham Pharmaceuticals, Old Powder Mills, Near Leigh, Tonbridge, Kent TN11 9AN (GB).
- (74) Agent: RUTTER, Keith; SmithKline Beecham, Corporate Intellectual Property, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).

(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published

With international search report.

(54) Title: HYDRATE OF 5-[4-[2- (N-METHYL-N- (2-PYRIDIL)AMINO) ETHOXY]BENZYL] THIAZOLIDINE-2, 4-DIONE MALEIC ACID SALT

(57) Abstract

A hydrate of 5-[4-[2- (N-methyl-N- (2-pyridyl)amino) ethoxy]benzyl] thiazolidine-2, 4-dione, maleic acid salt, characterised in that it: (i) comprises water in the range of from 0.2 to 1.1 % w/w; and (ii) provides an infrared spectrum containing peaks at 764 and 579 cm⁻¹; and/or (iii) provides an X-ray powder diffraction (XRPD) pattern substantially as set out in Figure II; a process for the preparation of such a compound, a pharmaceutical composition containing such a compound and the use of such a compound or composition in medicine.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
ΑU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
ΑZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
вв	Barbados	GH	Ghana	MG	Madagascar	ТJ	Tajikistan
BE	Belgium	GŇ	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
ВJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of Americ
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		•
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

HYDRATE OF 5-[4-[2-(N-METHYL-N-(2-PYRIDIL)AMINO)ETHOXY]BENZYL]THIAZOLIDINE-2,4-DIONE MALEIC ACID SALT

This invention relates to a novel pharmaceutical, to a process for the preparation of the pharmaceutical and to the use of the pharmaceutical in medicine.

International Patent Application, Publication Number WO94/05659 discloses certain thiazolidinedione derivatives having hypoglycaemic and hypolipidaemic activity including 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2.4-dione, maleic acid salt (hereinafter also referred to as "Compound (I)").

Compound (I) is disclosed solely as an anhydrous form. It has now been discovered that Compound (I) exists in a novel hydrated form which is particularly suitable for bulk preparation and handling. This can be prepared by an efficient, economic and reproducible process particularly suited to large scale preparation.

The novel hydrate also has useful pharmaceutical properties and in particular it is indicated to be useful for the treatment and/or prophylaxis of diabetes mellitus, conditions associated with diabetes mellitus and certain complications thereof.

Accordingly, the present invention provides a hydrate of 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione, maleic acid salt (the "Hydrate") characterised in that the Hydrate:

- 20 (i) comprises water in the range of from 0.2 to 1.1% w/w; and
 - (ii) provides an infra red spectrum containing peaks at 764 and 579 cm⁻¹; and/or
 - (iii) provides an X-ray powder diffraction (XRPD) pattern substantially as set out in Figure II.

Suitably, the water content of the Hydrate is in the range of from 0.4 to 0.9%w/w, especially 0.5 to 0.6% w/w, for example 0.54% w/w or 0.6 % w/w.

In one favoured aspect, the Hydrate provides an infra red spectrum substantially as set out in accordance with Figure I.

The Hydrate can exist in certain dehydrated forms which reversibly convert to the Hydrate when contacted with water, either in liquid or vapour form. The present invention encompasses all such reversibly rehydratable forms of the Hydrate.

The present invention encompasses the Hydrate isolated in pure form or when admixed with other materials, for example the known anhydrous form of Compound I, the above mentioned reversibly rehydratable forms or any other material.

Thus in one aspect there is provided the Hydrate in isolated form.

In a further aspect there is provided the Hydrate in pure form.

In yet a further aspect there is provided the Hydrate in crystalline form.

The invention also provides a process for preparing the Hydrate. characterised in that 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione,

5

10

15

25

30

maleic acid salt is crystallised from an aqueous alkanolic solvent, preferably containing from 2.0 to 2.5% by volume of water, most preferably 2.0 to 2.3%, for example 2.1 % by volume of water.

Suitable aqueous alkanolic solvents include aqueous ethanol, typically aqueous denatured ethanol, and aqueous isopropanol, or mixtures thereof.

Crystallisation and any recrystallisation is generally carried out at low to ambient temperature, such as in the range of between 0 to 30°C for example 25°C: alternatively crystallisation may be initiated at an elevated temperature, such as in the range of between 30°C and 60°C for example 35°C, and then completed by allowing the temperature of the solvent to cool to ambient or low temperature, such as in the range of between 0 to 30°C for example 25°C.

The crystallisation can be initiated by seeding with crystals of the Hydrate but this is not essential.

Compound I is prepared according to known procedures, such as those disclosed in WO94/05659. The disclosures of WO94/05659 are incorporated herein by reference.

When used herein the term 'prophylaxis of conditions associated with diabetes mellitus' includes the treatment of conditions such as insulin resistance, impaired glucose tolerance, hyperinsulinaemia and gestational diabetes.

Diabetes mellitus preferably means Type II diabetes mellitus.

Conditions associated with diabetes include hyperglycaemia and insulin resistance, especially acquired insulin resistance and obesity. Further conditions associated with diabetes include hypertension, cardiovascular disease, especially atherosclerosis, certain eating disorders, in particular the regulation of appetite and food intake in subjects suffering from disorders associated with under-eating, such as anorexia nervosa, and disorders associated with over-eating, such as obesity and anorexia bulimia. Additional conditions associated with diabetes include polycystic ovarian syndrome and steroid induced insulin resistance.

The complications of conditions associated with diabetes mellitus encompassed herein includes renal disease, especially renal disease associated with the development of Type II diabetes including diabetic nephropathy, glomerular sclerosis, nephrotic syndrome, hypertensive nephrosclerosis and end stage renal disease.

As used herein 'aqueous' with reference to a given solvent or solvent mixture refers to a solvent which contains sufficient water to provide Hydrate i.e having from 0.2 to 1.1 %w/w of water.

5

10

15

20

25

30

As mentioned above the compound of the invention has useful therapeutic properties: The present invention accordingly the Hydrate for use as an active therapeutic substance.

More particularly, the present invention provides the Hydrate for use in the treatment and/or prophylaxis of diabetes mellitus, conditions associated with diabetes mellitus and certain complications thereof.

The Hydrate may be administered <u>per se</u> or, preferably, as a pharmaceutical composition also comprising a pharmaceutically acceptable carrier. The formulation of the Hydrate and dosages thereof are generally as disclosed for Compound (I) in International Patent Application, Publication Number WO94/05659.

Accordingly, the present invention also provides a pharmaceutical composition comprising the Hydrate and a pharmaceutically acceptable carrier therefor.

The Hydrate is normally administered in unit dosage form.

The active compound may be administered by any suitable route but usually by the oral or parenteral routes. For such use, the compound will normally be employed in the form of a pharmaceutical composition in association with a pharmaceutical carrier, diluent and/or excipient, although the exact form of the composition will naturally depend on the mode of administration.

2.3

144

THE REP

Compositions are prepared by admixture and are suitably adapted for oral, parenteral or topical administration, and as such may be in the form of tablets, capsules, oral liquid preparations, powders, granules, lozenges, pastilles, reconstitutable powders, injectable and infusable solutions or suspensions, suppositories and transdermal devices. Orally administrable compositions are preferred, in particular shaped oral compositions, since they are more convenient for general use.

Tablets and capsules for oral administration are usually presented in a unit dose, and contain conventional excipients such as binding agents, fillers, diluents, tabletting agents, lubricants, disintegrants, colourants, flavourings, and wetting agents. The tablets may be coated according to well known methods in the art.

Suitable fillers for use include cellulose, mannitol, lactose and other similar agents. Suitable disintegrants include starch, polyvinylpyrrolidone and starch derivatives such as sodium starch glycollate. Suitable lubricants include, for example, magnesium stearate. Suitable pharmaceutically acceptable wetting agents include sodium lauryl sulphate.

Solid oral compositions may be prepared by conventional methods of blending, filling, tabletting or the like. Repeated blending operations may be used to

ì

5

10

15

20

25

30

distribute the active agent throughout those compositions employing large quantities of fillers. Such operations are, of course, conventional in the art.

Oral liquid preparations may be in the form of, for example, aqueous or oily suspensions, solutions, emulsions, syrups, or elixirs, or may be presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, for example sorbitol, syrup, methyl cellulose, gelatin, hydroxyethylcellulose, carboxymethyl cellulose, aluminium stearate gel or hydrogenated edible fats, emulsifying agents, for example lecithin, sorbitan monooleate, or acacia; non-aqueous vehicles (which may include edible oils), for example, almond oil, fractionated coconut oil, oily esters such as esters of glycerine, propylene glycol, or ethyl alcohol; preservatives, for example methyl or propyl p-hydroxybenzoate or sorbic acid, and if desired conventional flavouring or colouring agents.

For parenteral administration, fluid unit dose forms are prepared containing a compound of the present invention and a sterile vehicle. The compound, depending on the vehicle and the concentration, can be either suspended or dissolved. Parenteral solutions are normally prepared by dissolving the active compound in a vehicle and filter sterilising before filling into a suitable vial or ampoule and sealing. Advantageously, adjuvants such as a local anaesthetic, preservatives and buffering agents are also dissolved in the vehicle. To enhance the stability, the composition can be frozen after filling into the vial and the water removed under vacuum.

Parenteral suspensions are prepared in substantially the same manner except that the active compound is suspended in the vehicle instead of being dissolved and sterilised by exposure to ethylene oxide before suspending in the sterile vehicle. Advantageously, a surfactant or wetting agent is included in the composition to facilitate uniform distribution of the active compound.

In addition such compositions may contain further active agents such as anti-hypertensive agents and diuretics.

As is common practice, the compositions will usually be accompanied by written or printed directions for use in the medical treatment concerned.

As used herein the term 'pharmaceutically acceptable' embraces compounds, compositions and ingredients for both human and veterinary use: for example the term 'pharmaceutically acceptable salt' embraces a veterinarily acceptable salt.

The present invention further provides a method for the treatment and/or prophylaxis of diabetes mellitus, conditions associated with diabetes mellitus and certain complications thereof, in a human or non-human mammal which comprises administering an effective, non-toxic, amount of Hydrate to a human or non-human mammal in need thereof.

5

10

15

20

25

30

Conveniently, the active ingredient may be administered as a pharmaceutical composition hereinbefore defined, and this forms a particular aspect of the present invention.

In the treatment and/or prophylaxis of diabetes mellitus, conditions associated with diabetes mellitus and certain complications thereof Hydrate may be taken in doses, such as those described above.

Similar dosage regimens are suitable for the treatment and/or prophylaxis of non-human mammals.

In a further aspect the present invention provides the use of Hydrate for the manufacture of a medicament for the treatment and/or prophylaxis of diabetes mellitus, conditions associated with diabetes mellitus and certain complications thereof.

No adverse toxicological effects are indicated for the compounds of the invention in the above mentioned treatments.

The following examples illustrate the invention but do not limit it in any way.

٠,٠

5

Example 1: Preparation of hydrate of 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione, maleic acid salt.

5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2.4-dione free base (4.0 g) and maleic acid (1.40 g, 1.05 molar equivalents) were heated in denatured ethanol (40 ml) containing additional water (0.51 g, i.e. a total water content of approximately 2.1% (v/v)) to 60°C and held at this temperature for 30 minutes during which time a solution was obtained. The solution was filtered, re-heated to 50°C, and then cooled with stirring at a rate of 1 deg/min, resulting in crystallisation at 35°C.

- The resulting suspension was cooled to 25°C and stirred for two hours. The product was filtered, washed with 99% denatured ethanol (8 ml) and dried at 50°C in vacuo for 24 hours to give the title compound (4.38 g, 82%). The water content of the product was 0.54% w/w.
- 15 **CHARACTERISING DATA:** The following characterising data was generated for the Hydrate:

A Infrared

- The infrared absorption spectrum of a mineral oil dispersion of the Hydrate was obtained using a Nicolet 710 FT-IR spectrometer at 2 cm⁻¹ resolution. Data were digitised at 1 cm⁻¹ interval. The spectrum obtained is shown in Figure I. Peak positions are as follows: 3129, 2776, 1756, 1747, 1706, 1641, 1617, 1586, 1542, 1512, 1413, 1351, 1331, 1290, 1264, 1246, 1210, 1182, 1163, 1108, 1078, 1064, 1031, 1005, 075, 055, 036, 865, 764, 738, 710, 662, 619, 579, 557, 532, 535, and 50
- 25 1031, 1005, 975, 955, 926, 865, 764, 738, 719, 662, 619, 579, 557, 532, 525, and 508 cm⁻¹.

B X-Ray Powder Diffraction (XRPD)

The XRPD pattern of the Hydrate is shown below in Figure II and a summary of the XRPD angles and calculated lattice spacing characteristic of the Hydrate is given in Table I.

A PW1710 X-ray powder diffractometer (Cu X-ray source) was used to generate the spectrum using the following acquisition conditions:

Tube anode:

35

Cu

Generator tension:

40 kV

Generator current:

30 mA

Start angle:

3.5 °2θ 35.0 °2θ

40 End angle:

0.020

Time per step:

Step size:

4.550 s

Table I.

X-Ray Powder Diffraction Angles and Calculated Lattice Spacing Characteristic of the Hydrate.

Diffraction Angle	Lattice Spacing
(°2θ)	(Angstroms)
7.6	11.65
8.9	9.90
9.7	9.09
15.1	5.85
15.6	5.68
17.0	5.22
17.5	5.08
17.9	4.96
19.2	4.62
20.1	4.41
20.6	4.30
22.2	4.00
23.8	3.73
24.4	3.64
25.2	3.54
25.9	3.44
26.7	3.34
27.5	3.25
28.0	3.19
29.9	2.99
31.5	2.84

CLAIMS

- 1. A hydrate of 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione, maleic acid salt., characterised in that it:
 - (i) comprises water in the range of from 0.2 to 1.1% w/w; and
 - (ii) provides an infra red spectrum containing peaks at 764 and 579 cm⁻¹; and/or
 - (iii) provides an X-ray powder diffraction (XRPD) pattern substantially as set out in Figure II.

10

- 2. A hydrate according to claim 1, wherein the water content is in the range of from 05 to 0.6%w/w.
- 3. A hydrate according to claim 1 or claim 2, which provides an infra red spectrum substantially in accordance with Figure I.
 - 4. A hydrate according to any one of claims 1 to 3, which provides an X-ray powder diffraction (XRPD) pattern substantially as set out in Figure II.
- 20 5. A hydrate according to any one of claims 1 to 4, in isolated form.
 - 6. A hydrate according to any one of claims 1 to 5, in pure form.
 - 7. A hydrate according to any one of claims 1 to 6, in crystalline form.

25

- 8. A compound in the form of a rehydratable form of a hydrate according to any one of claims 1 to 7.
- 9. A process for preparing a hydrate according to claim 1, characterised in that 5 [4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione, maleic acid salt is crystallised from aqueous ethanol.
 - 10. A process according to claim 11, wherein the aqueous ethanol contains from 2% to 2.5% of water by volume.

35

11. A pharmaceutical composition comprising an effective, non-toxic amount of a hydrate according to claim 1 and a pharmaceutically acceptable carrier therefor.





5

10

- 12. A hydrate according to claim 1, for use as an active therapeutic substance.
- 13. A hydrate according to claim 1, for use in the treatment and/or prophylaxis of diabetes mellitus, conditions associated with diabetes mellitus and certain complications thereof.
- 14. The use of Hydrate for the manufacture of a medicament for the treatment and/or prophylaxis of diabetes mellitus, conditions associated with diabetes mellitus and certain complications thereof.
- 15. A method for the treatment and/or prophylaxis of diabetes mellitus, conditions associated with diabetes mellitus and certain complications thereof, in a human or non-human mammal which comprises administering an effective, non-toxic, amount of Hydrate to a human or non-human mammal in need thereof.







SUBSTITUTE SHEET (RULE 26)

			1 317 27 307	00104	
A. CLASSIF IPC 6	FICATION OF SUBJECT MATTER CO7D417/12 A61K31/425				
According to	International Patent Classification (IPC) or to both national classifica	ition and IPC			
B. FIELDS		on symbole)			
IPC 6	currentation searched (classification system followed by classification CO7D A61K	,,, aymio∪io)			
Documentati	ion searched other than minimum documentation to the extent that se	uch documents are incl	uded in the fields se	arched	
Electronic da	ata base consulted during the international search (name of data bas	se and. where practical	I, search terms used		
C. DOCUME	ENTS CONSIDERED TO BE RELEVANT		····	~ ~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	
Category 3	Citation of document, with indication, where appropriate, of the rele	evant passages		Relevant to claim No.	
X	WO 94 05659 A (SMITHKLINE BEECHAM; POOL COLIN RIPLEY (GB); ROMAN ROSHERWO) 17 March 1994 cited in the application see page 3, line 32; example 1			1-15	
X	EP 0 306 228 A (BEECHAM GROUP PLO 8 March 1989 see page 4, line 50; claim 1 			1-15	
Furti	ther documents are listed in the continuation of box C.	X Patent family	members are listed	in annex.	
considered to be of particular relevance "E" sartier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "X" document of particular relevance "Y" document of particular relevance "Y" document of particular relevance "Y" document of particular relevance "onvolve an inventive step where the considered novel ocument is combined with ments, such combination be in the art. "&" document member of the san			and not in conflict with and the principle or the cular relevance; the clered novel or canno ive step when the docular relevance; the clered to involve an inbined with one or michination being obvious and the control of the control	uffict with the application but sple or theory underlying the sple or theory underlying the sple or the considered to ear the document is taken alone sple an inventive step when the sple or more other such docurning obvious to a person skilled see patent family	
	2 March 1999	19/03/		·	
	mailing address of the ISA	Authorized officer			
rvanie and	mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040. Tx. 31 651 epo nl. Fax: (+31-70) 340-3016	Lauro,			

Form PCT/ISA/210 (second sheet) (July 1992)

information on patent family members

Inte onal Application No
PCT/EP 98/08154

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
WO 9405659	A	17-03-1994	AU	674880 B	16-01-1997
			AU	4973093 A	29-03-1994
			CA	2143849 A	17-03-1994
			CN	11 019 11 A,B	26-04-1995
			CN	1183275 A	03-06-1998
			CN	1183413 A	03-06-1998
			CN	1183276 A	03-06-1998
			CZ	9500565 A	15-11-1995
			EP	0658161 A	21-06-1995
			FI	951004 A	03-03-1995
			FI	982413 A	06-11-1998
			HU	72639 A	28-05-1996
			IL	106904 A	30-09-1997
÷			ĴΡ	2828777 B	25-11-1998
			JP	8501095 T	06-02-1996
			MX	9305397 A	31-01-1995
			NO	950852 A	03-03-1995
			NO	974646 A	03-03-1995
			NZ	255505 A	22-08-1997
			PL	307812 A	26-06-1995
			SG	48302 A	17-04-1998
			SI	9300452 A	30-06-1994
			SK	27795 A	09-08-1995
			US	5741803 A	21-04-1998
			ZA	9306509 A	16-06-1994
EP 0306228	 А	08-03-1989		2173888 A	09-03-1989
2. 0000220			CA	1328452 A	12-04-1994
			CA	1339902 A	09-06-1998
			DK	490288 A	05-03-1989
			EP	0842925 A	20-05-1998
			ĴΡ	10194970 A	28-07-1998
			JP	10194971 A	28-07-1998
			JР	1131169 A	24-05-1989
			ĴΡ	2614497 B	28-05-1997
			JР	2817840 B	30-10-1998
			ĴΡ	9183771 A	15-07-1997
			ĴΡ	2837139 B	14-12-1998
			ĴΡ	9183726 A	15-07-1997
			JP	9183772 A	15-07-1997
			PT	88410 A,B	31-07-1989
			ÜŚ	5646169 A	08-07-1997
			US	5002953 A	26-03-1991
				5521201 A	28-05-1996
			111		
			US		
			US	5232925 A	03-08-1993
			US US	5232925 A 5194443 A	03-08-1993 16-03-1993
			US	5232925 A	03-08-1993

Form PCT.TSA 210 (patent family annex) (July 1992)

This Page Blank (uspto)